

Further information

The Urology Clinic is located at the KEMH Centenary Clinic off Hensman Road in Subiaco. If travelling by car it is advisable to allow extra time to find a parking space.

It is not advisable to bring children to this appointment. Supervised care is available at the hospital crèche.

Telephone (08) 9340 1370 for more details.

Interpreters are available by appointment to assist you on the day of the test. Advise the appointment clerk when you are confirming your appointment if you require assistance from an interpreter.

Disclaimer: The advice and information contained herein is provided in good faith as a public service. However the accuracy of any statements made is not guaranteed and it is the responsibility of readers to make their own enquiries as to the accuracy, currency and appropriateness of any information or advice provided. Liability for any act or omission occurring in reliance on this document or for any loss, damage or injury occurring as a consequence of such act or omission is expressly disclaimed.

Produced by: Urogynaecology Clinic

Email: ogccu@health.wa.gov.au

Web site: <http://wnhs.health.wa.gov.au>

© July 2003 WNHS 0075 Rev 1

Reviewed August 2009

WOMEN AND NEWBORN HEALTH SERVICE

King Edward Memorial Hospital

374 Bagot Road Subiaco WA 6008

Telephone: (08) 9340 2222

Facsimile: (08) 9388 1780



Government of Western Australia
Department of Health



Delivering a Healthy WA

Urodynamic Studies

What is Urodynamics?

Urodynamics is a test that looks at the behaviour of the bladder. There are different types of urinary incontinence and they require different treatments.

This test will help ensure you have the best treatment for your bladder.

Patient information

Confirm your attendance at this appointment immediately by phoning (08) 9340 1355 between 8.30am and 3.00pm.

If you are unable to keep this appointment, phone at least 48 hours in advance of your appointment so we can allocate this time to another patient.

Stop taking any bladder therapy medications five days before your appointment as they may interfere with the accuracy of the test. These medications may include:

- Ditropan
- Proanthene
- Tofranil
- Vesicare
- Endep

You can continue to take this medication after completion of the Urodynamic Studies.

Note: If you are taking any of these medications for depression **do not stop** taking them. If you are unsure please contact your medical practitioner or the Urology Clinic.

If you have frequent urinary tract infections or feel you have symptoms of a urinary tract infection contact the Urology Nurse

Consultant on (08) 9340 2222 pager 3136, Monday to Friday 8.00am to 4.00pm to discuss.


Patient instructions

- Arrive for the appointment with a comfortably full bladder.
- One hour before your appointment drink two glasses of water, do not empty your bladder.
- You may eat and drink normally on the day of your test.
- Bring all of your medications to the appointment.
- You are attending this clinic because you have a bladder or prolapse problem. Do not feel embarrassed or upset if you have difficulty controlling your bladder during the test.

What happens during the test?

1. The test will take approximately 45 - 60 minutes to complete. It is not painful but you may find it a little uncomfortable.
2. You will be asked to empty your bladder in private into a special toilet.
3. A specimen of urine is collected to make sure you do not have a current bladder infection. A bladder infection may cause inaccurate results. If a bladder infection is suspected the test will be discontinued and another appointment made.
4. Fine tubes will be inserted into your bladder and vagina which will record pressure readings in your bladder throughout the test.

5. Your bladder will be filled with sterile water until you feel the need to pass urine.
6. You will be directed to cough at various stages during the test.
7. At the end of the test you will be allowed to empty your bladder again.



After the test

Occasionally some women develop a urinary tract infection after this test, so we advise you to increase your fluid intake for 24 hours after the test. You will be given instructions for bladder care. If you are concerned that you have an infection, please see your own local doctor.