



NEONATAL Medication Monograph

# NORADRENALINE (Norepinephrine)

This document should be read in conjunction with this [DISCLAIMER](#)





**Restricted:** Requires Neonatologist or Cardiologist review within 24 hours of initiation

**⚠ HIGH RISK Medication**

<b>Presentation</b>	Vial: 4mg/4mL = 1000microgram/mL
<b>Description</b>	Noradrenaline stimulates beta <sub>1</sub> -adrenergic and alpha-adrenergic receptors. This causes increased contractility and heart rate as well as vasoconstriction to increase systemic blood pressure and coronary blood flow.
<b>Indications</b>	Vasopressor activity for vasodilatory shock, e. g septic shock <i>Note: Usually used in conjunction with dopamine to treat profound hypotension, in the setting of sepsis when other inotropes have failed.</i>
<b>Contraindications</b>	Hypotension from hypovolaemia except as an emergency measure to maintain cerebral and coronary perfusion until volume replaced
<b>Precautions</b>	Ensure adequate circulatory volume to minimise need for vasoconstrictors Avoid hypertension: close monitoring of BP and adjustment if needed Use with extreme caution in patients with profound hypoxia or hypercarbia may produce ventricular tachycardia or fibrillation use with extreme caution Flows <2mL/hr may cause swings in blood pressure
<b>Dosage</b>	<b><i>Dose stated as noradrenaline base</i></b>  <b>IV Infusion :</b> <i>Initial Dose</i> 0.05 – 0.1 microgram/ kg/ minute  <i>Titrate dose in small increment as below :</i> 0.05 – 0.5 microgram/ kg/ minute Higher doses up to a <b>maximum of 2microgam/ kg/ minute</b> ; have been used as per neonatologist or cardiologist

<b>Dosage Adjustment</b>	Titrate initial dose in small increments as required
<b>Adverse Reactions</b>	<b>Common:</b> anxiety, palpitations, headache, extravasation which may cause local tissue necrosis.
	<b>Serious:</b> hypertension which may be associated with bradycardia. Prolonged use of high doses can result in renal impairment and reduced peripheral tissue perfusion.
<b>Interactions</b>	Benzyloxyphenol , ganciclovir, indomethacin, insulin, sodium bicarbonate, phenobarbitone Check Y-site compatibility with Pharmacy
<b>Compatible Fluids</b>	Glucose 5%, Sodium Chloride 0.9%
<b>Preparation</b>	IV available from CIVAS (KEMH+PCH)  <b>Infusion Calculation:</b> <b>Dilute</b> 300 microgram (0.3mL ) of noradrenaline base per kilogram of baby's weight and dilute to 50mL with compatible diluent  <b>Concentration = 1mL/hr = 0.1 microgram/kg/minute</b>  <b>E.g</b> To prepare an infusion for a 780g infant Weight = 0.78kg Dose in Infusion = 300 microgram x 0.78 = 234 microgram Dilute 234 microgram (≈0.23mL) to 50mL with compatible fluid.
<b>Administration</b>	Infuse into a central venous catheter using a rate-controlled infusion device
<b>Monitoring</b>	Continuous blood pressure monitoring Monitor IV site for extravasation
<b>Storage</b>	Store at room temperature, below 25°C Protect ampoules from light.
<b>Notes</b>	<ul style="list-style-type: none"> <li>Do not cease infusion abruptly</li> <li>Correct hypovolaemia before commencing noradrenaline</li> </ul>
<b>Related clinical guidelines</b>	<a href="#">Congenital Diaphragmatic Hernia (CHD)</a> <a href="#">Cardiac: Routine Post-Operative Care</a>

<b>References</b>	<p>Takemoto CK, Hodding JH, Kraus DM. Pediatric &amp; neonatal dosage handbook with international trade names index: a universal resource for clinicians treating pediatric and neonatal patients. 23rd ed. Hudson (Ohio): Lexicomp; 2017. 2,401 p.1445</p> <p>Truven Health Analytics. Norepinephrine. In: NeoFax [Internet]. Greenwood Village (CO): Truven Health Analytics; 2020 [cited 2020 Apr 06]. Available from: <a href="https://neofax.micromedexsolutions.com/">https://neofax.micromedexsolutions.com/</a></p> <p>Australian Medicines Handbook. Children’s Dosing Companion. Noradrenaline (norepinephrine) In: Australian Medicines Handbook Children’s Dosing Companion [Internet]. Adelaide (South Australia): Australian Medicines Handbook Children’s Dosing Companion; 2019 [cited 2020 Apr 06].</p> <p>Shann F. Drug doses. Seventeenth ed. Parkville (Victoria): Collective P/L; 2017. 75</p> <p>Society of Hospital Pharmacists of Australia. Noradrenaline (Norepinephrine). In: Australian Injectable Drugs Handbook [Internet]. [St Leonards, New South Wales]: Health Communication Network; 2020 [cited 2020 Apr 06]. Available from: <a href="http://aidh.hcn.com.au">http://aidh.hcn.com.au</a></p>
-------------------	---

Keywords:	Norepinephrine (noradrenaline)		
Publishing:	<input checked="" type="checkbox"/> Intranet	<input checked="" type="checkbox"/> Internet	
Document owner:	Head of Department - Neonatology		
Author / Reviewer:	KEMH & PCH Pharmacy / Neonatology Directorate		
Date first issued:	March 2008	Version:	3.0
Last reviewed:	April 2020	Next review date:	April 2023
Endorsed by:	Neonatal Directorate Management Group	Date:	April 2020
Standards Applicable:	NSQHS Standards: 1  Governance, 3  Infection Control, 4  Medication Safety, 8  Acute Deterioration		

**Printed or personally saved electronic copies of this document are considered uncontrolled.**

**Access the current version from the WNHS website.**

**For any enquiries relating to this guideline, please email [KEMH.PharmacyAdmin@health.wa.gov.au](mailto:KEMH.PharmacyAdmin@health.wa.gov.au)**

© Department of Health Western Australia 2019