



ADULT MEDICATION GUIDELINE

Oxycodone

Scope (Staff): All WNHS Staff

Scope (Area): Obstetrics and Gynaecology

This document should be read in conjunction with the [Disclaimer](#).

Quick Links

[Dose](#)

[Administration](#)

[Monitoring](#)

[Pregnancy and Breastfeeding](#)

Restrictions

[Formulary: Restricted](#)

[HIGH RISK Medication](#) 

Medication Class

Analgesics

Presentation

Immediate Release Tablets: 5mg

Immediate Release Capsules: 10mg, 20mg

Oral liquid: 1mg/mL

Controlled Release Tablets: 10mg, 15mg, 20mg, 30mg, 40mg, 80mg

Oxycodone/Naloxone (Targin®) Combined Controlled Release Tablets:

2.5/1.25mg, 5/2.5mg, 10/5mg, 20/10mg, 40/20mg

Storage

Store at room temperature, below 25°C

Schedule 8 Medication

Dose

Moderate to Severe Pain

Doses vary and depend on the indication and previous analgesic requirements.

Titrate dose according to response, respiratory rate and sedation score.

Suggested starting dose (immediate release): 5mg every 2 to 4 hours

Suggested starting dose (controlled release tablets): total daily dose as determined for immediate release products; give up to half total daily dose every 12 hours

Administration

Controlled release tablets: Swallow whole. Do not break, crush or chew.

Oxycontin®: After putting the tablet in the mouth, make sure to wash it down immediately with an adequate amount of water. Review If patient have problems swallowing the tablets.

Monitoring

Respiratory rate, sedation score

Pregnancy

Considered safe to use (lowest effective dose for the shortest duration possible)

For more information, please contact [KEMH Obstetric Medicines Information Service](#).

Breastfeeding

Considered safe to use during breastfeeding at the lowest effective doses for the shortest duration possible.

Dose should not be higher than 30 mg a day for no longer than 3 days. Observe the breastfed infant for adverse effects such as excessive drowsiness, poor feeding or sleeping pattern changes.

For more information, please contact [KEMH Obstetric Medicines Information Service](#).

Related Policies, Procedures & Guidelines

Department of Health

[Recommendations for prescribing analgesia on discharge following surgery or acute injury](#)

[CCC Report on the Supply and Management of Schedule 8 Controlled Drugs at Certain Public Hospitals in Western Australia](#)

WNHS Clinical Practice Guidelines:

[Palliative Care](#)

[Acute Pain Service](#)

Patient Information Leaflet:

[Medicines used to manage pain](#)

[Post-operative pain control](#)

WNHS Pharmaceutical and Medicines Management Guidelines:

[Anaesthetic technician competency framework for handling of controlled substances](#)

[Reporting of S8 or S4R medicine discrepancies](#)

References

Australian Medicines Handbook. Oxycodone. In: Australian Medicines Handbook [Internet]. Adelaide (South Australia): Australian Medicines Handbook; 2021 [cited 2021 May 7]. Available from: <https://amhonline.amh.net.au/>









The Royal Women's Hospital. Oxycodone. In: Pregnancy and Breastfeeding Medicines Guide [Internet]. Parkville (Victoria): The Royal Women's Hospital; 2021 [cited 2021 Jul 30]. Available from: <https://thewomenspbmg.org.au/>

APILAM. Oxycodone. In: e-lactancia.org. APILAM: association for promotion of and cultural and scientific research into breastfeeding; 2002 updated 10 jun 2014; accessed 30 jul 2021. Available from <http://e-lactancia.org/breastfeeding/oxycodone/product/>

Hale TW. Oxycodone. In: HalesMeds.com [Internet]. New York: Springer Publishing Company; c2021 [updated 2019 Oct 4; cited 2021 Jul 30]. Available from: <https://www.halesmeds.com/>

Therapeutic Guidelines. Oral drugs for severe, acute nociceptive pain in adults. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2020 Dec [cited 2021 Jul 30]. Available from: <https://tgldcdp-tg-org-au.kelibresources.health.wa.gov.au/etgAccess>

Therapeutic Guidelines. Oral drugs for moderate, acute nociceptive pain in adults. In: eTG complete [Internet]. West Melbourne (Victoria): Therapeutic Guidelines; 2020 Dec [cited 2021 Jul 30]. Available from: <https://tgldcdp-tg-org-au.kelibresources.health.wa.gov.au/etgAccess>

Keywords	Oxycodone, oxycontin, targin, naloxone, opioid, analgesia, analgesic, schedule 8, s8, controlled drug, controlled medicine, pain				
Document Owner:	Chief Pharmacist				
Author/ Reviewer	KEMH Pharmacy Department				
Version Info:	4.0				
Date First Issued:	07/2015	Last Reviewed:	30/07/2021	Review Date:	30/07/2024
Endorsed by:	Medicines and Therapeutics Committee			Date:	16/09/2021
NSQHS Standards Applicable:	<input checked="" type="checkbox"/>  Std 1: Clinical Governance <input type="checkbox"/>  Std 2: Partnering with Consumers <input type="checkbox"/>  Std 3: Preventing and Controlling Healthcare Associated Infection <input checked="" type="checkbox"/>  Std 4: Medication Safety		<input type="checkbox"/>  Std 5: Comprehensive Care <input type="checkbox"/>  Std 6: Communicating for Safety <input type="checkbox"/>  Std 7: Blood Management <input type="checkbox"/>  Std 8: Recognising and Responding to Acute Deterioration		
Printed or personally saved electronic copies of this document are considered uncontrolled. Access the current version from WNHS HealthPoint.					

The health impact upon Aboriginal people has been considered, and where relevant incorporated and appropriately addressed in the development of this document (insert ISD Number). (Please refer to the Aboriginal Health Impact Statement and Declaration for Department of Health and Health Service Provider Guidelines – please delete once you have completed this).

This document can be made available in alternative formats on request for a person with a disability.

© Women and Newborn Health Service 2021

Copyright of this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.