



Government of **Western Australia**
North Metropolitan Health Service
Women and Newborn Health Service



Antenatal expression of colostrum for the vulnerable baby



Breastfeeding provides many emotional and physical benefits for both you and your baby.

A Baby Friendly Health Initiative Accredited Hospital (BFHI), this initiative aims to reduce practices in maternity settings that have a negative impact on breastfeeding.

It is recommended to breast feed exclusively for six months.

Antenatal colostrum expression is a strategy specific to avoiding the use of formula supplementation for vulnerable babies. These include babies born to a diabetic mother; babies born less than 39 weeks gestation; babies who are smaller or larger than expected for gestational age; and babies weighing less than 2500g at birth.

Providing colostrum to infants at risk of hypoglycaemia has been shown to improve exclusive breastfeeding and decrease formula supplementation [Tozier, 2013].

Colostrum

- Colostrum is the first milk that is produced by the breast during pregnancy and the first few days after the birth of your baby.
- Colostrum contains many protective substances for the immune system of the newborn baby.
- Colostrum is low in volume but high in energy and helps with the passage of meconium (the baby's first bowel motion) which assists in preventing jaundice (yellow skin colour).

If you are considering expressing colostrum antenatally, please discuss this with your doctor, midwife, diabetes educator or lactation consultant.

It is important you do not express antenatally if:

- You have a history of threatened or actual preterm labour
- You have cervical incompetence
- You have a cervical suture preventing preterm labour
- You have had any bleeding in your pregnancy
- You have placenta praevia

Commencing expressing in the antenatal period for vulnerable babies

- Hand expressing can be started from 36 weeks gestation
- Start with three to five minutes of hand expressing on each breast twice per day
- Cease hand expressing immediately if you are having tightening or contractions

Hand expressing must be pain free. If you require additional assistance or support, please contact the Breast Feeding Centre of WA on 6458 1844

How do I express by hand?



- Wash your hands with soap and water and dry on a clean towel prior to expressing.
- Use a clean container, medicine cup or syringe.
- Ensure you are sitting comfortably.
- Gently stroke the breast towards the nipple. This stimulates the reflex.
- Place the fingers underneath the breast, so the first finger is just below and the thumb is just above the areola about 3-4cm back from the nipple.
- Gently **press** the fingers and thumb pads (not fingertips) back towards the chest.
- Then **compress** the breast tissue and hold briefly.
- **Release** the breast tissue.
- Do not squeeze or pinch your nipple.
- Repeat the action in a rhythm similar to baby's sucking.
- Rotate the position of the fingers and thumb around the breast to express all of the breast.
- When colostrum is pearling or dripping easily, it is time to collect the colostrum.
- Express both breasts in turn while the colostrum is dripping.
- Express both breasts twice during each expressing session. The expressed milk is stored in the freezer during your pregnancy.

What do I do with my expressed colostrum?

Colostrum can be collected twice each day in the same syringe. You will need to store the syringe in the fridge between expressions.

At the end of the collecting day, the colostrum can be frozen – place the capped syringe into the freezer.

Label the syringe with a sticker, detailing the date and time of the expression.

Your colostrum will remain frozen until you come to hospital. Make sure each container is labelled with your name, hospital record number, date and the time the colostrum was expressed.

Freshly expressed colostrum should be cooled before being added to previously expressed chilled or frozen colostrum.

Storage of your expressed colostrum:

Breast milk	Room temperature	Refrigerator	Freezer
Freshly expressed into a clean container	6 to 8 hours (26°C or lower)	3 to 5 days (4°C or lower)	Two weeks in freezer compartment inside a refrigerator
	Store in refrigerator if one is available	Store in back of refrigerator where it is coldest	Three months in freezer section of refrigerator (with separate door)
			6 to 12 months in deep freeze (-18°C or lower)

* Australian Dietary Guidelines for Children and Adolescents table. Copyright Commonwealth of Australia reproduced by permission.

When you come to hospital for the birth of your baby:

- Bring your frozen expressed colostrum clearly labeled into the hospital and discuss storage with your midwife.

What happens after the birth of my baby?

- Providing you and baby are well, you will be encouraged to have skin-to-skin contact with your baby in the first one to two hours following birth.
- Breastfeeding is encouraged and should be unrestricted.
- If your baby is experiencing difficulties or is not interested in breastfeeding at this time, you will be encouraged to hand express some colostrum to give to your baby. You may also use any colostrum you have brought into hospital with you.
- Should you and your baby continue to experience breastfeeding challenges or blood glucose levels are low, the midwife and paediatrician will guide you on further treatment and feeding.

There is wide variation in the amount of colostrum women can express antenatally. There is no evidence to suggest that a woman who expresses small amounts of colostrum antenatally is at risk of low breast milk supply.

References

1. Cox (2010). An ethical dilemma: should recommending antenatal expressing and storage of colostrum continue? *Breastfeeding Review* 18(3) 5-7
2. Forster et al (2011). Diabetes and antenatal milk expressing: A pilot project to inform development of a randomised control trial. *Midwifery* 27(2)209-214
3. National Health and Medical Research Council 2012, *Infant feeding guidelines: Summary*. Canberra, ACT: National Health and Medical Research Council.
4. Forster DA, Moorhead AM, Jacobs SE, Davis PG, Walker SP, McEgan KM, Opie GF, Donath SM, Gold L, McNamara C, Aylward A, Ford R, Amir LH 2017, Advising women with diabetes in pregnancy to express breastmilk in late pregnancy (Diabetes and Antenatal Milk Expressing [DAME]): a multicentre, unblinded, randomised controlled trial. *The Lancet*, 389(10085): 2204-2213.
5. Tozier, P. K. (2013). Colostrum versus formula supplementation for glucose stabilisation in newborns of diabetic mothers. *Journal of Obstetric, Gynaecology and Neonatal Nursing*. 42,619-628
6. Singh et al (2009) effect of antenatal expression of Breast milk at Term in Reducing Breast Feeding Failures. *MAIFI* 65:131-133



Women and Newborn Health Service

King Edward Memorial Hospital
374 Bagot Road, Subiaco WA 6008

☎ (08) 6458 2222

🌐 kemh.health.wa.gov.au

Compiled by: OGID

Produced by: Women and Newborn Health Service - Revised Feb 2022

This document can be made available
in alternative formats on request.