



Laparoscopy

Patient discharge information

Information following your operation

You have just had a Laparoscopy. Complications after this operation are rare, however the following advice is provided to help you avoid complications and to recognise them if they occur.

1. Do not try to do too much. Rest when you get home. You should be able to resume normal activities after a couple of days, although the effects of the anaesthetic may make you feel tired.
2. You will have some discomfort in the places where the incisions (cuts) were made in your abdomen (tummy). You may also have muscular aches and pains from the anaesthetic and pain in your shoulders from the gas put into your abdomen during the operation. These discomforts will settle in a couple of days.
3. You may take Panadol and non-steroidal/anti-inflammatories (e.g. Ibuprofen) or prescribed tablets for pain, cramps or headaches.
4. You may use pads for any bleeding.
5. You may shower as usual.
6. If you also had a curette (scraping of the womb) a small amount of bleeding, lasting 10 to 14 days, is common. The bleeding may be heavier with increased activity, such as lifting.

For one week you should not:

- swim, bathe or take a spa
 - have sexual intercourse
 - use tampons.
7. After a curette, your next period will usually begin 3 to 6 weeks following your operation. If you did NOT have a curette, your next period should arrive at the usual time.
 8. If you are taking the oral contraceptive pill, your next period will begin as usual at the end of the packet.
 9. If your period has not arrived after 6 weeks, please check with your doctor (GP).
 10. An infection may be present if you have:
 - heavy bleeding, i.e. needing to change your pad every 10 to 20 minutes
 - fever (temperature, chills)
 - pain in the lower part of your abdomen not relieved by pain medication
 - an offensive (smelly) discharge from the vagina
 - red and painful wounds on your abdomen.

Heavy bleeding and infection may be serious. If you think you have one or both of these complications, please contact your GP the Emergency Centre at KEMH on (08) 6458 1433.

Following a general anaesthetic

1. It is essential that someone stays overnight with you.
2. You must not drive a car or operate machinery for at least 24 hours after your operation.
3. You must not consume alcohol for 24 hours.
4. You should not make any important decisions or sign any important papers for at least 24 hours after your operation.
5. You should avoid any activity that requires the use of skill or judgement, for example delicate work, cooking etc. for 12 hours.
6. You should avoid social engagements for 12 hours.

Follow-up care

It is important to visit your GP or the KEMH clinic for a post-operative check up.

Operation performed

Follow-up appointment

Special instructions



Women and Newborn Health Service

King Edward Memorial Hospital
374 Bagot Road, Subiaco WA 6008
Telephone: (08) 6458 2222
kemh.health.wa.gov.au

Produced by Obstetrics and Gynaecology Clinical Care Unit

© North Metropolitan Health Service 2020

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair dealing for the purposes of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without written permission of the State of Western Australia.



This document can be made available
in alternative formats on request.