




# Syphilis testing in pregnancy

Patient information



## Acknowledgement of Country

We acknowledge the Aboriginal people of the many traditional lands and language groups of Western Australia and pay respect to their elders past and present. North Metropolitan Health Service recognises, respects and values Aboriginal cultures as we walk a new path together.



# Contents

Pregnant or planning pregnancy? ..... 4

What is syphilis?..... 5

What is congenital syphilis? ..... 6

Syphilis testing ..... 7

If you do not have syphilis ..... 8

If you have syphilis..... 9

Syphilis treatment ..... 10

# Pregnant or planning pregnancy?

Get tested for syphilis as soon as you know you are pregnant.



# What is syphilis?

**Syphilis** is a sexually transmitted infection. There is currently a syphilis outbreak in WA.



Common symptoms of syphilis are shown (left), but some people have no symptoms (right).

## What is congenital syphilis?


When a pregnant woman has a syphilis infection, syphilis can pass to her unborn baby. This can cause miscarriage, premature birth and harm to the baby, occasionally resulting in the baby's death. Treatment can prevent this. This is why testing is so important.



# Syphilis testing

All pregnant women should be tested for syphilis **at least** three times during pregnancy:

1. At first doctor's visit
2. At 28 weeks
3. At 36 weeks.



## If you do **not** have syphilis

**If your test shows you do not have syphilis,** you still need to be tested regularly during your pregnancy. If symptoms (see page 5) develop, tell your doctor/midwife and have extra testing.





## If you have syphilis


**If your test shows you have syphilis**, you will need treatment and have more frequent testing. Your doctor or midwife will explain this.



# If you have syphilis


## Syphilis treatment

Antibiotics are used to treat syphilis and can prevent the baby becoming infected.



## If you have syphilis

Do not have any sexual activity until seven days after you and your partner/s have been treated.



Your doctor or midwife will arrange follow-up testing to make sure the antibiotics have worked.

## If you have syphilis

You will be asked about who you have had sex with. It is very important that they get a blood test and treatment.

This will stop you getting reinfected with syphilis and protect your baby.

If you think you have been reinfected with syphilis, speak to your doctor or midwife as soon as you can.




# If you have syphilis

The management will include:



Antibiotics



No sex for seven days after treatment



Testing and treating partner/s



Extra syphilis testing

# If you have syphilis

You will be looked after by a team, such as your GP, hospital midwives and doctors, and public health nurses who might call to check how you're going. We work together to look after you and your pregnancy.



GP



Public health  
nurses



Hospital midwives  
and doctors

## If you have syphilis

After you have your baby, your healthcare team will check on you and your baby before you go home and may arrange follow-up appointments.





## Further information

Visit [syphilis in pregnancy \(healthywa.wa.gov.au\)](https://healthywa.wa.gov.au).

or scan the QR code



### Boorloo (Perth) Public Health Unit

 3/311 Wellington Street Perth WA 6000

 1300 623 292

 (08) 9222 8599

This document can be made available  
in alternative formats on request.

© North Metropolitan Health Service 2024



We are proud to be a smoke-free site.  
Thank you for not smoking or vaping  
in any buildings or on our grounds.